Functional Groups

Dr. Sapna Gupta

Classes of Compounds

- Classification based on **functional group**.
- The functional group is where the reaction usually takes place.
- Alkanes don't have a functional group and are the least reactive of all classes.
- Three broad classes
 - Hydrocarbons
 - Compounds containing oxygen
 - Compounds containing nitrogen.
- Organic compounds are classified also as aliphatic (straight chain) and aromatic (benzene like).
- Functional groups can be classified in "families" or homologous series. These series increase by 1 carbon to give the next compound.

Nomenclature - IUPAC

- Prefix tells the number of carbon atoms.
- Suffix tells the name of the functional group.


	Number of	Number of		
Prefix	c arbo ns	Prefix	carbons	
meth-	1	unde c-	11	
eth-	2	do de c-	12	
prop-	3	tride c-	13	
but-	4	tetrade c-	14	
pe nt-	5	pe ntade c	- 15	
he x-	6	he xade c-	16	
he pt-	7	he ptade c	- 17	
oct-	8	octadec-	18	
no n-	9	no nade c-	19	
de c-	10	eicos-	20	

Example:

Methane: meth – 1 carbon; ane – alkane Butanol: but – 4 carbons; ol – alcohol Ethanal: eth – 2 carbons; al – aldehyde Propanone: prop – 2 carbons; one - ketone


				Family			
	Alkane	Alkene	Alkyne	Aromatic	Haloalkane	Alcohol	Ether
Functional group	C—H and C—C bonds	C=C	—C≡C—	Aromatic ring	C ∷:	—с—ён 	cё
General formula	RH	$\begin{array}{l} \text{RCH} = \text{CH}_2 \\ \text{RCH} = \text{CHR} \\ \text{R}_2\text{C} = \text{CHR} \\ \text{R}_2\text{C} = \text{CR}_2 \end{array}$	RC≡CH RC≡CR	ArH	RX	ROH	ROR
Specific example	CH ₃ CH ₃	CH ₂ =CH ₂	HC≡CH	\bigcirc	CH ₃ CH ₂ Cl	CH ₃ CH ₂ OH	CH ₃ OCH ₃
IUPAC name	Ethane	Ethene	Ethyne	Benzene	Chloroethane	Ethanol	Methoxymethane
Common name ^a	Ethane	Ethylene	Acetylene	Benzene	Ethyl chloride	Ethyl alcohol	Dimethyl ether

^aThese names are also accepted by the IUPAC.



Solved Example

• Circle and name all the functional groups in the following compound.



Key Words/Concepts

- Fundamentals of naming organic compounds
- Functional Groups